

Fermionic systems with charge correlations

FERDINANDO MANCINI

*Dipartimento di Fisica "E.R. Caianiello" - Unità di ricerca INFN di Salerno
Università degli Studi di Salerno, I-84081 Baronissi (SA), Italy*

PACS. 71.10.-w – Theories and models of many-electron systems.

PACS. 71.10.Fd – Lattice fermion models (Hubbard model, etc.).

PACS. 71.27.+a – Strongly correlated electron systems; heavy fermions.

Abstract. – In this paper, we show that a system of localized particles, satisfying the Fermi statistics and subject to finite-range interactions, can be exactly solved in any dimension. In fact, in this case it is always possible to find a finite closed set of eigenoperators of the Hamiltonian. Then, the hierarchy of the equations of motion for the Green's functions eventually closes and exact expressions for them are obtained in terms of a finite number of parameters. For example, the method is applied to the two-state model (equivalent to the spin-1/2 Ising model) and to the three-state model (equivalent to the extended Hubbard model in the ionic limit or to the spin-1 Ising model). The models are exactly solved for any dimension d of the lattice. The parameters are self-consistently determined in the case of $d = 1$.

How do we study systems characterized by the presence of a huge number of interacting particles? The standard procedure is a perturbative one. At first, the interaction is completely ignored or approximately taken into account as an external field (mean field). Then, a perturbative expansion is considered. By construction, this scheme works well only when the interaction is sufficiently weak, but miserably fails when the correlations among the particles are quite strong. This is the case for all new highly interacting materials [Fulde(1995)]. An alternative procedure has been suggested in Refs. [Mancini and Avella(2003), Mancini(2003)], where the main ingredient is the use of composite operators. In the presence of interactions the properties of the original (bare) particles are modified and new particles (quasi-particles) should be considered. The difficulty with this approach [named composite operator method (COM)] is that the number of composite operators rapidly increases with the number of degrees of freedom and one must again resort to approximate treatments [Mancini and Avella(2004)]. Recently, I have found that there is a large class of systems for which it is always possible to find a finite closed set of eigenoperators of the Hamiltonian and exact solution can be obtained. The purpose of this letter is to define this class, and to illustrate through some examples the technique of calculations. Details of calculations and further results will be given elsewhere.

We consider a system of q species of particles, satisfying Fermi statistics, subject to finite-range interactions, localized on the sites of a Bravais lattice. We suppose that the mass of the particles is very large and/or the interaction is so strong that the kinetic energy is negligible and the particles are frozen on the lattice sites. For a two-body interaction the Hamiltonian

takes the form

$$H = \sum_{\mathbf{i}a} V_a n_a(i) + \frac{1}{2} \sum_{\mathbf{i}j} \sum_{ab} V_{ab}(\mathbf{i}, \mathbf{j}) n_a(i) n_b(j) \quad (1)$$

where $\mathbf{i}$ stands for the lattice vector $\mathbf{R}_i$ and $i = (\mathbf{i}, t)$. $c_a(i)$ and $c_a^\dagger(i)$ are annihilation and creation operators of particles of species a , in the Heisenberg picture, satisfying canonical anti-commutation relations.

$$\begin{aligned} \{c_a(\mathbf{i}, t), c_b^\dagger(\mathbf{j}, t)\} &= \delta_{ab} \delta_{\mathbf{ij}} \\ \{c_a(\mathbf{i}, t), c_b(\mathbf{j}, t)\} &= \{c_a^\dagger(\mathbf{i}, t), c_b^\dagger(\mathbf{j}, t)\} = 0 \end{aligned} \quad (2)$$

V_a represents an external field acting on the particles a . $n_a(i) = c_a^\dagger(i) c_a(i)$ is the charge density operator of the fields a . $V_{ab}(\mathbf{i}, \mathbf{j})$ describes the two-body interactions. It is immediate to see that the $n_a(i)$ satisfies the equation of motion $i \frac{\partial n_a(i)}{\partial t} = [n_a(i), H] = 0$. Then, in order to use the equations of motion and Green's function (GF) formalism we must start from the Heisenberg equation for the a species

$$i \frac{\partial c_a(i)}{\partial t} = V_a c_a(i) + \sum_{j\mathbf{b}} V_{ab}(\mathbf{i}, \mathbf{j}) n_b(j) c_a(i) \quad (3)$$

where, without loss of generality, we put $V_{aa}(\mathbf{i}, \mathbf{i}) = 0$. We see that the dynamics has generated another field operator $\varphi_a(i) = \sum_{j\mathbf{b}} V_{ab}(\mathbf{i}, \mathbf{j}) n_b(j) c_a(i)$ of higher complexity. By taking time derivatives of increasing order, more and more complex operators are generated. These operators are composite operators [Mancini and Avella(2003), Mancini and Avella(2004)], as they are all expressed in terms of the original fields $c_a(i)$ and $c_a^\dagger(i)$. Now, I assert that: **because of the algebra satisfied by the charge density operator $n_a(i)$, the hierarchy of composite operators closes.** The proof of the statement is the following. For the sake of simplicity, let us take $V_{ab}(\mathbf{i}, \mathbf{j}) = 2d\delta_{ab}V\alpha_{\mathbf{ij}}$, where d is the dimensionality of the system, $\alpha_{\mathbf{ij}}$ is the projector on the nearest-neighbor sites and V is the strength of the interaction. I am considering systems with first-nearest neighbor interactions, but the following proof can be extended to systems with second, third, $\dots$ -nearest neighbor interactions. By increasing the range of interactions the number of eigenoperators will increase. The time derivative of order k is

$$\left(i \frac{\partial}{\partial t}\right)^k c_a(i) = V_a^k c_a(i) + \sum_{m=1}^k m V_a^{k-m} (2dV)^m [n^\alpha(i)]^m c_a(i)$$

where $n^\alpha(i) = \sum_{j\mathbf{b}} \alpha_{\mathbf{ij}} n_b(j)$. Because of (2), the operator $[n^\alpha(i)]^m$ obeys the following recurrence rule

$$[n^\alpha(i)]^m = \sum_{p=1}^{2qd} A_p^{(m)} [n^\alpha(i)]^p \quad \sum_{p=1}^{2qd} A_p^{(m)} = 1 \quad (4)$$

The coefficients $A_p^{(m)}$ are rational numbers that can be easily determined by the algebra and the structure of the lattice. Then, for $k > 2qd$ no additional composite operators are generated and the equations of motion close. By putting $n = 2qd + 1$, we construct a n -multiplet composite operator

$$\psi(i) = \begin{pmatrix} \psi_1(i) \\ \psi_2(i) \\ \vdots \\ \psi_n(i) \end{pmatrix} \quad (5)$$

which satisfies the Heisenberg equation

$$i\frac{\partial\psi(i)}{\partial t} = [\psi(i), H] = \epsilon\psi(i) \quad (6)$$

where the $n \times n$ matrix ϵ will be denominated as the energy matrix. Note that we are using a vectorial notation: the field $\psi_m(i)$ is itself a multiplet of rank q . Once the composite operator $\psi(i)$ and the energy matrix have been determined, an exact solution of the Hamiltonian can be obtained. Let us define the retarded Green's function

$$G(i, j) = \theta(t_i - t_j) \langle \{\psi(i), \psi^\dagger(j)\} \rangle \quad (7)$$

where $\langle \dots \rangle$ denotes the quantum-statistical average over the grand canonical ensemble. By means of the Heisenberg equation (6) we obtain in momentum space the equation

$$[\omega - \epsilon]G(\mathbf{k}, \omega) = I(\mathbf{k}) \quad (8)$$

where $I(\mathbf{k})$ is the Fourier transform of the normalization matrix: $I(\mathbf{i}, \mathbf{j}) = \langle \{\psi(\mathbf{i}, t), \psi^\dagger(\mathbf{j}, t)\} \rangle$. The solution of Eq. (8) is

$$G(\mathbf{k}, \omega) = \sum_{m=1}^n \frac{\sigma^{(m)}(\mathbf{k})}{\omega - E_m + i\delta} \quad (9)$$

where E_m are the eigenvalues of the energy matrix ϵ . The spectral density matrices $\sigma^{(m)}(\mathbf{k})$ are calculated by means of the formula [Mancini and Avella(2003)]

$$\sigma_{ab}^{(m)}(\mathbf{k}) = \Omega_{am} \sum_{c=1}^n \Omega_{am}^{-1} I_{cb}(\mathbf{k}) \quad (10)$$

where Ω is the $n \times n$ matrix whose columns are the eigenvectors of the matrix ϵ . The spectral density matrices $\sigma^{(m)}(\mathbf{k})$ satisfy the sum rule $\sum_{m=1}^n E_m^p \sigma^{(m)}(\mathbf{k}) = M^{(p)}(\mathbf{k})$, where $M^{(p)}(\mathbf{k})$ are the spectral moments defined as

$$M^{(p)}(\mathbf{k}) = F.T. \langle \{ (i\partial/\partial t)^p \psi(\mathbf{i}, t), \psi^\dagger(\mathbf{j}, t) \} \rangle \quad (11)$$

$F.T.$ stays for the Fourier transform. The correlation function (CF) $C(i, j) = \langle \psi(i)\psi^\dagger(j) \rangle$ can be immediately calculated from (9) and one obtains

$$C(\mathbf{k}, \omega) = \pi \sum_{m=1}^n \delta[\omega - E_m] T_m \sigma^{(m)}(\mathbf{k}) \quad (12)$$

with $T_m = 1 + \tanh(\beta E_m/2)$ and $\beta = 1/k_B T$ is the inverse temperature.

Equations (9) and (12) are an exact solution of the model Hamiltonian (1). However, the knowledge of the GF is not fully achieved yet. The algebra of the field $\psi(i)$ is not canonical: as a consequence, the normalization matrix $I(\mathbf{k})$ in the equation of motion (8) contains some unknown static correlation functions. These correlators are expectation values of operators not belonging to the chosen basis $\psi(i)$, and should be self-consistently calculated.

According to the scheme of calculations proposed by COM [Mancini and Avella(2003), Mancini(2003), Mancini and Avella(2004)], one way of calculating the unknown correlators is by specifying the representation where the GF is realized. The procedure is the following. From the algebra it is possible to derive several relations among the operators. We will call algebra constraints (AC) all possible relations among the operators dictated by the algebra.

This set of relations valid at microscopic level must be satisfied also at macroscopic level (i.e., when expectations values are considered). Use of these considerations leads to some self-consistent equations which will be used to fix the unknown correlator appearing in the normalization matrix. An immediate set of rules is given by the equation

$$\langle \psi(i) \psi^\dagger(i) \rangle = \frac{1}{N} \sum_{\mathbf{k}} \frac{1}{2\pi} \int_{-\infty}^{+\infty} d\omega C(\mathbf{k}, \omega) \quad (13)$$

where the l.h.s. is fixed by the AC and the boundary conditions compatible with the phase under investigation, while in the r.h.s. the correlation function $\langle \psi(i) \psi^\dagger(i) \rangle$ is computed by means of (12).

Another set of AC can be derived by observing that there exist some operators, O , which project out of the Hamiltonian a reduced part $OH = OH_0$. When H_0 and $H_I = H - H_0$ commute, the quantum statistical averages of O over the complete Hamiltonian must coincide with the average over the reduced Hamiltonian H_0

$$Tr\{Oe^{-\beta H}\} = Tr\{Oe^{-\beta H_0}\} \quad (14)$$

Another relation comes from the requirement of time translational invariance which leads to the condition that the spectral moments, defined by Eq. (11), must satisfy the following relation

$$M_{nm}^{(p)}(\mathbf{k}) = M_{mn}^{(p)}(\mathbf{k})^* \quad (15)$$

It can be shown that if (15) is violated, states with negative norm appear in the Hilbert space. Of course, the above rules are not meant to be exhaustive and, in principle, more conditions might be needed.

We now apply the above procedure to two specific models. It goes without saying that the proposed models belong to the class of the Potts models [Potts(1952)]. We consider a d -dimensional cubic Bravais lattice.

The two-state model

As first example, we consider only one species of particles. By taking $V_a = -\mu$, $V_{ab}(\mathbf{i}, \mathbf{j}) = 2dV\delta_{ab}\alpha_{\mathbf{i}, \mathbf{j}}$ the Hamiltonian (1) takes the form

$$H = -\mu \sum_{\mathbf{i}} v(i) + Vd \sum_{\mathbf{i}} v(i)v^\alpha(i) \quad (16)$$

where hereafter for a generic operator $\Phi(i)$ we use the notation $\Phi^\alpha(\mathbf{i}, t) = \sum_{\mathbf{j}} \alpha_{\mathbf{i}, \mathbf{j}} \Phi(\mathbf{j}, t)$. μ is the chemical potential and we put $v(i) = c^\dagger(i)c(i)$; we are ignoring the subindex a . By means of the transformation $v(i) = \frac{1}{2}[1 + S(i)]$ the Hamiltonian (16) can be cast in the form

$$H = E_0 - h \sum_{\mathbf{i}} S(i) - dJ \sum_{\mathbf{i}} S(i)S^\alpha(i) \quad (17)$$

where $E_0 = (-\frac{1}{2}\mu + \frac{1}{4}dV)N$, $h = \frac{1}{2}(\mu - dV)$, $J = -\frac{1}{4}dV$. Hamiltonian (17) is just the d -dimensional spin-1/2 Ising model [Ising(1925)] with nearest neighbor interactions in presence of an uniform external magnetic field.

To solve the Hamiltonian (16) let us consider the composite operator

$$\psi(i) = \begin{pmatrix} \psi_1(i) \\ \psi_2(i) \\ \vdots \\ \psi_{2d+1}(i) \end{pmatrix} = \begin{pmatrix} c(i) \\ v^\alpha(i)c(i) \\ \vdots \\ [v^\alpha(i)]^{2d}c(i) \end{pmatrix} \quad (18)$$

This field is an eigenoperator of the Hamiltonian (16)

$$i\frac{\partial}{\partial t}\psi(i) = [\psi(i), H] = \epsilon\psi(i) \quad (19)$$

The energy matrix ϵ has the rank $(2d+1) \times (2d+1)$ and can be calculated by means of (3) and the recurrence rule (4). Proof of the relation (4) and the explicit expressions of $A_m^{(p)}$ for this model will be given elsewhere. The eigenvalues of the energy matrix are given by

$$E_m = -\mu + (m-1)V \quad \{m = 1, 2, \dots, (2d+1)\} \quad (20)$$

The retarded GF $G(i, j)$ and the CF $C(i, j)$ can be exactly calculated by applying the scheme of calculations illustrated above. By using the anti-commutation rules (2) and the symmetry relations (15), we obtain $I(\mathbf{i}, \mathbf{j}) = \delta_{\mathbf{i}, \mathbf{j}} I_0$, where the matrix I_0 depends on the parameters $\kappa^{(p)} = \langle [v^\alpha(i)]^p \rangle$ ($p = 1, \dots, 2d$). Then, the matrices $\sigma^{(m)}$ are calculated by means of (10). We have an exact solution of the Ising model for dimensions $d = 1, 2, 3$, where all the properties of the model are expressed in terms of the self-consistent parameters $\kappa^{(p)}$. To have quantitative results we must calculate $\kappa^{(p)}$ in terms of the external parameters μ, T and V . We now recall the AC (13). Because of the relation (4) there are only $2d+1$ independent elements of the matrix $C = \langle \psi(i)\psi^\dagger(i) \rangle$ and we obtain the self-consistent equations

$$\kappa^{(p)} - \lambda^{(p)} = \frac{1}{2} \sum_{m=1}^{2d+1} T_m \sigma_{1,p}^{(n)} \quad (p = 1, \dots, 2d+1) \quad (21)$$

where $T_m = 1 + \tanh(\beta E_m/2)$ and $\sigma^{(n)}$ are the spectral density matrices. New correlation functions $\lambda^{(p)} = \langle v(i)[v^\alpha(i)]^p \rangle$ appear and the set of self-consistent equations (21) is not sufficient to determine all unknown parameters. One needs more conditions. In the case $d = 1$, these extra conditions can be obtained by using property (14). By means of the algebraic relation $c^\dagger(i)v(i) = 0$, we have $c^\dagger(i)e^{-\beta H} = c^\dagger(i)e^{-\beta H_0}$, where $H_0 = H - 2Vv(i)v^\alpha(i)$. By requiring that (14) be satisfied, we can derive the relation $\frac{C_{1k}}{C_{11}} = \frac{C_{1k}^{(0)}}{C_{11}^{(0)}}$ where $C_{1k} = \langle c(i)c^\dagger(i)[v^\alpha(i)]^{k-1} \rangle$ and $C_{1k}^{(0)} = \langle c(i)c^\dagger(i)[v^\alpha(i)]^{k-1} \rangle_0$. Here $\langle \dots \rangle_0$ denotes the thermal average with respect to H_0 . By exploiting these relations we can arrive to the following equation

$$C_{13} = \frac{1}{2}C_{12} \left(1 + \frac{C_{12}}{C_{11}} \right) \quad (22)$$

This equation closes the system of self-consistent equations and we have a complete solution of the model. Higher-order correlation functions can be studied by similar techniques. All details of the calculations will be given in a separate work, where we will show that all the well-known results [Baxter(1982), Goldenfel(1992), Lavis and Bell(1999)] of the one-dimensional Ising model are recovered.

Three-state model

We consider now a second example. Let us consider two species of particles, say a and b , and take $V_a = V_b = -\mu$, $V_{ab}(\mathbf{i}, \mathbf{j}) = U\delta_{\mathbf{i}, \mathbf{j}}\delta_{ab} + 2dV\alpha_{\mathbf{i}, \mathbf{j}}$. Then, the Hamiltonian (1) becomes

$$H = \sum_{\mathbf{i}} [-\mu n(i) + UD(i) + Vn(i)n^\alpha(i)] \quad (23)$$

where $n(i) = n_a(i) + n_b(i)$ and $D(i) = n_a(i)n_b(i) = \frac{1}{2}n(i)[n(i) - 1]$ are the total particle density and double occupancy operators, respectively. This Hamiltonian is just the extended

Hubbard model in the ionic limit, where U and V are the on-site and inter-site Coulomb interaction, respectively. The two species of particles, a and b , are in this case electrons with spin up and down, respectively. By means of the transformation $n(i) = [1 + S(i)]$, (23) can be cast in the form

$$H = -dJ \sum_{\mathbf{i}} S(i) S^\alpha(i) + \Delta \sum_{\mathbf{i}} S^2(i) - h \sum_{\mathbf{i}} S(i) + E_0 \quad (24)$$

where $E_0 = (-\mu + dV)N$, $h = \mu - 2dV - \frac{1}{2}U$, $J = -dV$, $\Delta = \frac{1}{2}U$. Hamiltonian (24) is just the Ising spin-1 model [Blume et al.(1971)] with nearest-neighbor interactions in presence of a crystal field Δ and an external magnetic field h .

To solve the Hamiltonian (23) let us consider the composite operators

$$\psi^{(x)}(i) = \begin{pmatrix} \psi_1^{(x)}(i) \\ \psi_2^{(x)}(i) \\ \vdots \\ \psi_{4d+1}^{(x)}(i) \end{pmatrix} = \begin{pmatrix} x(i) \\ x(i)[n^\alpha(i)] \\ \vdots \\ x(i)[n^\alpha(i)]^{4d} \end{pmatrix} \quad (25)$$

where $x = \xi, \eta$, being $\xi(i) = [1 - n(i)]c(i)$ and $\eta(i) = n(i)c(i)$ the Hubbard operators. Note that $c(i)$, $\xi(i)$, $\eta(i)$ are doublet operators with components a and b . By means of (3) and (4), these fields are eigenoperators of the Hamiltonian (23)

$$\begin{aligned} i \frac{\partial}{\partial t} \psi^{(\xi)}(i) &= [\psi^{(\xi)}(i), H] = \epsilon^{(\xi)} \psi^{(\xi)}(i) \\ i \frac{\partial}{\partial t} \psi^{(\eta)}(i) &= [\psi^{(\eta)}(i), H] = \epsilon^{(\eta)} \psi^{(\eta)}(i) \end{aligned} \quad (26)$$

where $\epsilon^{(\xi)}$ and $\epsilon^{(\eta)}$ are the energy matrices, of rank $(4d+1) \times (4d+1)$, which can be calculated by means of the equations of motion (26) and the recurrence relation (4). The eigenvalues $E_m^{(\xi)}$ and $E_m^{(\eta)}$ of the energy matrices are

$$\begin{aligned} E_m^{(\xi)} &= -\mu + (m-1)V \\ E_m^{(\eta)} &= -\mu + U + (m-1)V \end{aligned} \quad \{m = 1, 2, \dots, (4d+1)\}$$

The retarded GF $G^{(xy)}(i, j) = \langle R[\psi^{(x)}(i) \psi^{(y)\dagger}(j)] \rangle$ and the CF $C^{(xy)}(i, j) = \langle \psi^{(x)}(i) \psi^{(y)\dagger}(j) \rangle$, with $x, y = \xi, \eta$, can be exactly calculated by applying the scheme of calculations illustrated above. By using the anti-commutation relations (3) and the symmetry relations (15), straightforward calculations show that $I^{(\xi\eta)}(\mathbf{i}, \mathbf{j}) = I^{(\eta\xi)}(\mathbf{i}, \mathbf{j}) = 0$, while

$$\begin{aligned} I_{1,k}^{(\xi\xi)}(\mathbf{i}, \mathbf{j}) &= \delta_{\mathbf{ij}} [\kappa^{(k-1)} - \lambda^{(k-1)}] \\ I_{1,k}^{(\eta\eta)}(\mathbf{i}, \mathbf{j}) &= \delta_{\mathbf{ij}} \lambda^{(k-1)} \end{aligned} \quad (k = 1, \dots, 4d+1)$$

where $\kappa^{(p)} = \langle [n^\alpha(i)]^p \rangle$, $\lambda^{(p)} = \frac{1}{2} \langle n(i) [n^\alpha(i)]^p \rangle$. The CF $C^{(xx)}(i, j)$, ($x = \xi, \eta$), have the expressions

$$C^{(xx)}(i, j) = \delta_{\mathbf{ij}} \frac{1}{2} \sum_{m=1}^{4d+1} T_m^{(x)} \sigma^{(xm)} e^{-iE_m^{(x)}(t_i - t_j)} \quad (27)$$

where $T_m^{(x)} = 1 + \tanh(\beta E_m^{(x)}/2)$. The spectral density matrices $\sigma^{(xm)}$ are expressed in terms of the elements $I_{1k}^{(xx)}$. We have an exact solution of the model for dimensions $d = 1, 2, 3$,

where all the properties are expressed in terms of the self-consistent parameters $\kappa^{(p)}$ and $\lambda^{(p)}$. In order to determine the correlators $\kappa^{(p)}$ and $\lambda^{(p)}$, we use the AC constraints (13). By means of the algebraic relations $\xi_c(i)\xi_c^\dagger(i) + \eta_c(i)\eta_c^\dagger(i) = 1 - n_c(i)$, with $c = a, b$, we obtain for the paramagnetic phase the following self-consistent equations

$$\kappa^{(k-1)} - \lambda^{(k-1)} = C_{1k}^{(\xi\xi)} + C_{1k}^{(\eta\eta)} \quad (k = 1, \dots, 4d + 1) \quad (28)$$

The number of equations is not sufficient to fix all the parameters and more equations are needed. In the case of $d = 1$ these extra conditions can be obtained by using the property (14). By means of the algebraic relations

$$\begin{aligned} \xi^\dagger(i)n(i) &= 0 & D^p(i) &= D(i) \\ \xi^\dagger(i)D(i) &= 0 & D(i)n^p(i) &= 2^p D(i) \end{aligned} \quad (29)$$

we have

$$\begin{aligned} \xi^\dagger(i)e^{-\beta H} &= \xi^\dagger(i)e^{-\beta H_0} \\ D(i)e^{-\beta H} &= D(i) \sum_{p=0}^4 h_p [n^\alpha(i)]^p e^{-\beta H_0} \end{aligned} \quad (30)$$

where $H_0 = H - 2Vn(i)n^\alpha(i)$, and the h_p 's are known functions of the potential V . By requiring that (14) be satisfied we can derive a set of three self-consistent equations, which added to (28) allow us to determine all 8 parameters $\kappa^{(p)}$ and $\lambda^{(p)}$ ($p = 1, \dots, 4$). Calculations and results will be presented elsewhere.

Summarizing, we have shown that a system of localized particles, satisfying Fermi statistics, subject to finite-range interactions, can be described in terms of a closed set of eigenoperators. For the case of nearest neighbor interactions, the number of these composite operators is equal to $(2qd + 1)$, where q is the number of species of particles and $d = 1, 2, 3$ is the dimensionality of the system. The Green's functions and the correlation functions can be exactly calculated and are expressed in terms of a set of self-consistent parameters. For example, we have considered two models: the two-state model (equivalent to the spin- $\frac{1}{2}$ Ising model) and the three-state model (equivalent to the extended ionic Hubbard model and to the spin-1 Ising model). For these models the parameters are calculated in the case of $d = 1$. For higher dimensions more self-consistent equations are needed. This problem is now under investigation.

* * *

The author is grateful to Dr. A. Avella for stimulating discussions, a very friendly collaboration and his careful reading of the manuscript.

REFERENCES

- [Fulde(1995)] P. Fulde, *Electron Correlations in Molecules and Solids* (Springer-Verlag, Berlin Heidelberg New York, 1995), 3rd ed.
- [Mancini and Avella(2003)] F. Mancini and A. Avella, Eur. Phys. J. B **37**, 37 (2003).
- [Mancini(2003)] F. Mancini, in *Highlights in Condensed Matter Physics*, edited by A. Avella, R. Citro, C. Noce, and M. Salerno (American Institute of Physics, New York, 2003), pp. 240–257.
- [Mancini and Avella(2004)] F. Mancini and A. Avella, Advances in Physics **53**, 537 (2004).
- [Potts(1952)] R. Potts, Proc. Camb. Phil. Soc. **48**, 106 (1952).
- [Ising(1925)] E. Ising, Z. Physik **31**, 253 (1925).

- [Baxter(1982)] R. Baxter, *Exactly solved models in statistical mechanics* (Academic Press, London, 1982).
- [Goldenfel(1992)] N. Goldenfel, *Lectures on phase transitions and the renormalization group* (Perseus Publishing, Reading, Massachusetts, 1992).
- [Lavis and Bell(1999)] D. Lavis and G. Bell, *Statistical Mechanics of Lattice Systems*, vol. 1 of *Text and Monographs in Physics* (Springer-Verlag, Berlin, 1999), 2nd ed.
- [Blume et al.(1971)] M. Blume, V. Emery, and R. Griffiths, Phys. Rev. A **4**, 1071 (1971).